

研究業績

<分光学に関する研究>

1) アミロース・デンプン等多糖類のヨウ素呈色反応による物理化学的研究

半田隆、相川秀夫、石井忠浩、折笠令吉、鮎川彌一、工業化学雑誌、72、527 (1969)

2) Molecular Weight- and Solvent-effect on Magnetic Circular Dichroism of Polystyrenes in Solution.

T. Ishii, Y. Utena, T. Handa, Rept, Prog, Polym, Phys, Japan, 16, 87 (1976)

3) ポリスチレンおよびポリ-p-アルキルスチレン溶液のエキシマー発光

石井忠浩、松下秀鶴、半田隆、高分子論文集、32、211 (1975)

4) Effect of Tacticity on Excimer Formation of Poly(p-methylstyrenes) in Solution,

T. Ishii, S. Matsunaga, T. Handa, Makromol. Chem., 177, 283 (1976)

5) Fluorescence Quenching of Poly(N-vinylcarbazole) in Solution by Organic Acid.

T. Ishii, Y. Utena, T. Handa, Rept, Prog, Polym, Phys, Japan, 19, 399 (1976)

6) Exciplex Fluorescence Spectra of Copolymer Between Styrene and 2-vinylpyridine in Solution in the Presence of Organic Acid.

T. Ishii, Y. Utena, T. Handa, Rept, Prog, Polym, Phys, Japan, 19, 403 (1973)

7) フェニルカルバミン酸メチルの光転位反応

高仲善明、山縣知之、石井忠浩、半田隆、日本化学会誌、462、(1976)

8) A Study of Mechanism of Excimer Formation through Fluorescence Quenching for Isotactic and Atactic Polystyrenes and Poly(p-methylstyrenes in Solution.)

T. Ishii, T. Handa, S. Matsunaga Macromol. Chem., 178 2351 (1977)

9) Mechanism of Excimer Formation of Rigid and Flexible Polymers.

T. Ishii, Y. Utena, T. Handa, S. Mori, K. Takagi, Rept, Prog, Polym, Phys, Japan, 20, 415 (1977)

10) Exciplex Fluorescence Spectra of Poly-2-vinylpyridine in 1,2-Dichloroethane in the Presence of Trifluoroacetic Acid in Solution.

T. Ishii, Y. Utena, T. Handa, Rept, Prog, Polym, Phys, Japan, 20, 411 (1977)

11) Molecular Weight Effect on Polymer Motion of Polystyrene in the Complex Formation of Phenyl Rings with Acceptors.

T. Hand, T. Osawa, T. Ishii, Rept, Prog, Polym, Phys, Japan, 20, 423 (1977)

13) The Reaction of Poly-2-Vinylpyridine with Molecular Iodine.

(I) The Effect of Donor Concentration in Polymer.

T. Hand, T. Osawa, T. Ishii, S. Goto Rept, Prog, Polym, Phys, Japan, 20, 427 (1977)

14) The Reaction of Poly-2-Vinylpyridine with Molecular Iodine.

(II) The Effect of Temperature.

T. Hand, T. Osawa, T. Ishii, S. Goto Rept, Prog, Polym, Phys, Japan, 20, 431 (1977)

15) ポリウレタンの紫外線に対する挙動

山縣、石井、高仲、半田、日本化学会誌、1496 (1977)

16) The Effect of Molecular Weight on Excimer Formation of Polystyrenes in Solution.

T. Ishii, T. Handa, S. Matsunaga Macromolecules, 11, 41 (1978)

- 17)ポリウレタン樹脂の光反応における一重項酸素の関与
山縣、石井、高仲、半田、 高分子論文集 35, 349 (1978)
- 18)Emission Spectra of Poly(4-vinylpyridine) in Solution in the Presence of Trifluoroacetic Acid.
T. Ishii, T. Handa, Y. Utena, T. Moribe Rept, Prog, Polym, Phys, Japan, 21, 357 (1978)
- 19)Solvent Effect on Rate Constants of Excimer Formation and Dissociation of Flexible and Rigid
Vinylaromatic Polymers in Solution.
T. Ishii, T. Handa, S. Mori, Y. Utena, Rept, Prog, Polym, Phys, Japan, 21, 361 (1978)
- 20)The Evaluation of the Rate Constant of Excimer Formation of Poly(N-vinylcarbazole)
in Solution.
T. Ishii, T. Handa, Y. Utena, S. Mori, Rept, Prog, Polym, Phys, Japan, 21, 365 (1978)
- 21)The Reaction of Poly-2-vinylpyridine with Molecular Iodine (III).
T. Handa, T. Osawa, K. Suzuki, T. Ishii, S. Goto, Rept, Prog, Polym, Phys, Japan, 21, 369 (1978)
- 22)Steric Effects on Excimer Formation of Isotactic and Atactic Polystyrene and
Poly(*para*-methylstyrenes) in Solution.
T. Ishii, T. Handa, S. Matsunaga, J. Polym. Sci. Part A-2, 17, 811 (1979)
- 23)Molecular Weight Dependence of Magnetic Circular Dichroism of Polystyrenes in Solution.
T. Ishii, Y. Utena, T. Handa Makromol. Chem., 180, 175 (1979)
- 24)ポリウレタン樹脂の光反応における一重項酸素の影響と安定剤の効果
山縣、石井、高仲、半田、日本化学会誌、 1079 (1979)
- 25)4-メチル-*m*-フェニレン=ジイソシアナートから合成されたジウレタンの光反応
山縣、石井、高仲、半田、日本化学会誌、 1085 (1979)
- 26)ポリウレタン塗料の光劣化におよぼす酸化防止剤の効果
山縣、石井、高仲、半田、 高分子論文集、36, 767 (1979)
- 27)Photodegradation of Polyurethane Model Compound.
T. Ishii, T. Handa, T. Yamagata, S. Mori, K. Muto, Rept, Prog, Polym, Phys, Japan, 22, 547 (1979)
- 28)On the Blue Color of Triiodide Ions in Starch and Starch Fractions. I. Characterization of the
Changes in Absorption and Circular Dichroism Spectra of Triiodide Ions in Amylose with the DP.
T. Handa, H. Yajima, Biopolymers, 18, 873 (1979)
- 29)On the Blue Color of Triiodide Ions in Starch and Starch Fractions. II. Characterization of the
Changes in Absorption and Circular Dichroism Spectra of Triiodide Ions in Amylose with the DP.
T. Handa, H. Yajima, Biopolymers, 19, 723 (1980)
- 30)On the Blue Color of Triiodide Ions in Starch and Starch Fractions. III. Characterization of the
Changes in Absorption and Circular Dichroism Spectra of Triiodide Ions in Amylose with the DP.
T. Handa, H. Yajima, T. Kajiura, Biopolymers, 19, 1723 (1980)
- 31)Spectroscopic Properties of Homooligopeptides of L-Tryptophan in Ethanol.
T. Ishii, T. Handa, K. Muto, M. Ueki, S. Ikeda,
Reports on Progress in Polymer Physics in Japan, 24, 571-579 (1981)

- 32) Emission Properties of (2-Pyridyl)-(CH₂)_n-(2-Pyridyl) in Solution in the Presence of Trifluoroacetic Acid.
T. Ishii, T. Handa, Y. Utena, T. Yamauchi
Reports on Progress in Polymer Physics in Japan, 24, 449-452 (1981)
- 33) Enthalpy Change of Amylose Triiodide Complex Formation.
T. Handa, H. Yajima, T. Yamamura, T. Ishii, H. Aikawa, Starke, 32, 194 (1980)
- 34) Conformation of Amylose-Iodine-Iodide Complex in Aqueous Solution.
T. Handa, H. Yajima, Biopolymers, 20, 2051 (1981)
- 35) Deep Blueing Mechanism of Triiodide Ions in Amylose Being Associated with Its Conformation.
T. Handa, H. Yajima, T. Ishii, T. Nishimura, Acc Symposium Series 150, "Solution Properties of Polysaccharides", Ed. David A. Brant, Chapter 31, P455 (1981)
- 36) On the Purple Color of Triiodide Ions in Amylopectin.
H. Yajima, T. Handa, T. Nishimura, H. Aikawa, Rep. Prog. Polym. Phys. Jpn. 24, 669 (1981)
- 37) Fluorescence Spectra of Tryptophan Residues in Human Insoluble Protein.
Handa, Sato, Ishii, Kato, Rep. Prog. Polym. Phys. Jpn. 25, 719 (1982)
- 38) Effect of the Solution State of Amylose on the Coloring and Conformation of Its I₃⁻ Complex
H. Yajima, T. Handa, T. Ono, S. Mayama, H. Aikawa Rep. Prog. Polym. Phys. Jpn. 25, 693 (1982)
- 39) Effect of the Solution State of Amylose on the Coloring and Conformation of Its I₃⁻ Complex II.
Evaluation of the Occupied Ratios of the Being Species as a Function of Aging Time.
H. Yajima, T. Handa, T. Ono, S. Mayama, H. Aikawa Rep. Prog. Polym. Phys. Jpn. 26, 693 (1983)
- 40) エーテルの赤外多光子解離反応
真嶋、荒井、石井、豊田、半田、レーザー科学研究、5、134、(1983)
- 41) ピリジン系化合物のレーザー光分解の研究(I) 4-ビニールピリジンのKrFレーザーによる分解
松沼、石井、半田、荒井
- 42) "KrF Laser" Laser Photolysis and radiolysis of 4-Vinylpyridine
S. Matsunaga, T. Ishii, T. Handa, S. Arai
Chemistry Letters, 1121-1124 (1984)
- 43) Excimer Emission in Protonated Pyridine System. II. Excimer Emission of Protonated Dipyritylalkanes in Solution at Room Temperature and 77° K
T. Handa, Y. Utena, H. Yajima, Phys. Chem. 88, 5150 (1984)
- 44) Excimer Emission in Protonated Pyridine Systems. 1. Fluorescence Spectroscopy of Protonated Pyridine System and Its Methyl Derivatives in Rigid Glass Solution at 77° K.
T. Handa, Y. Utena, H. Yajima, T. Ishii, H. Morita, Phys. Chem. 90, 2589 (1986)
- 45) Excimer Emission in Protonated Pyridine Systems. 3. Excimer Emissions of Poly-2-&-4-Vinylpyridines in Solutions in the Presence of Trifluoroacetic Acid.
T. Handa, Y. Utena, H. Yajima, R. Katayama, T. Ishii, T. Yamauchi, Phys. Chem. (in Press)
- 46) Excimer Emission in Protonated Pyridine Systems. 4. Effect of Molecular Weight on Dimer-like Excimer Formation of Poly-2-Vinylpyridine in Rigid Glass Solution in the presence of Trifluoroacetic Acid at 77° K.
Y. Utena, H. Yajima, T. Ishii, T. Handa, Europ. Polym. J. (in Press)

47)Circular Dichroism of Backbone and Side Chains Separated from Natural Circular Dichroism of Poly-L-Tryptophan by Fluorescence Detected Circular Dichroism Method.

K. Muto, H. Mochizuki, R. Yoshida, T. Ishii, T. Handa, J. Am. Chem. Soc. (in Press)